#### The first characterisation of meat consumption across sociodemographic, lifestyle and anthropometric groups in Switzerland: Th Life Sciences und Facility Management menuCH 💙 LGI Institut für Leber results from the National Nutrition Survey menuCH

Universität Zürich<sup>™</sup>

**ETH** zürich



Autnors: Linda Tschanz<sup>1</sup>, Ivo Kaelin<sup>2</sup>, Anna Wróbel<sup>2</sup>, Giulia Pestoni<sup>3</sup>, Sabine Rohrmann<sup>3</sup>, Isabelle Herter-Aeberli<sup>1</sup>, Janice Sych<sup>4</sup> <sup>1</sup> Human Nutrition Laboratory, Department of Health Sciences and Technology, ETH Zurich, Switzerland <sup>2</sup> Institute of Applied Simulation, ZHAW School of Life Sciences and Facility Management, Schloss 1,8820 Waedenswil, Switzerland <sup>3</sup> Division of Chronic Disease Epidemiology, Epidemiology, Biostatistics and Prevention Institute, University of Zurich, Hirschengraben 84, 8001 Zurich, Switzerland <sup>4</sup> Institute of Food and Beverage Innovation, ZHAW School of Life Sciences and Facility Management, Einsiedlerstrasse 34, 8820 Waedenswil, Switzerland

## Introduction

4

## What types of meat consume the residents of Switzerland?

- Based on annual sales data, consumption of all meat categories except white meat is decreasing<sup>1</sup>.
- According to the Swiss Health Survey 2017, 53% of the population consumes meat more frequently than the recommended 3 times per week, however no statements can be made about compliance with the recommended consumption amount of 100 - 120 g/day.
- Consumption amounts were recently assessed by two 24-houers dietary recall interviews in the National Nutrition Survey menuCH representative for adults (18 – 75 years) living in the three main References: 1 Nahrunasmittelbilanz 2013 = Bilan alimentaire 2013. Agristat. 2015. language regions of Switzerland. 2 Schweizerische Gesundheitsbefragung 2017 : Übersicht. BFS, 2018.

## Method using the menuCH data

**Ouantification** of all, processed, red and white meat consumption was performed applying weights for sex, age, major area of Switzerland, marital status, household size nationality, season and weekday. Multiple linear regression was conducted on energy standardised and weighted consumption data (g/1,000 kcal) across sociodemographic, lifestyle and anthropometric variables for all, processed, red and white meat with the total population and sex specific. All regressions were adjusted for sex, language region, age, BMI, education, household status, income, physical activity, smoking, health status and dieting for weight loss. Significant results p-value < 0.05 in the table are presented in bold.

Quantification of meat consumption		Meat category consumption [g/day]												
		То	tal _	109	138	78	106	116	117	124	117	104	91	
37 g/day	The menuCH me category consum % of the po All meat Processed meat Red meat White meat	eat- ers pulation 89%		2.5 42.7 36.9 26.8	Sex 3.5 55.2 48.2		Languag 3.1 45.1 83.1 24.3	1 6 36.8 45.3 82.2	0.5 39.3 44.7 32.2	3.5 50 32 38.2	Age gri 3.2 42.8 41.4 29.9	2-5 40.5 37 23.9	0.8 39.1 34.5	
43 g/day	27 g/day		1	All OH	Men v	sh Meat	German	Frend	ttalian Meat	Age 18-20 Red Meat	- r	ge 45-59 pe ps	2	

# **Regression Results**

# Multiple linear regression [g/1000 kcal]

			I Populaiton n = 2057	Me	en n = 933	Women n = 1124		
		Coe		Coeff.	95% CI	Coeff.	95% CI	
			Vleat	000111	0070 01	000111	0070 01	
Sex								
	Men	0.0	ref.					
	Women	-10.1	[-13.6 ; -6.6]					
angu	age regions							
	German	0.0	ref.	0.0	ref.	0.0	ref.	
	French	7.0	[3.1 ; 10.8]	2.7	[-3.5; 8.9]	10.1	[5.4 ; 14.8	
	Italian	11.9	[4.7 ; 19.1]	11.8	[0.6; 23.1]	11.5	[2.5 ; 20.5	
Age-g	roups							
	18-29 years	7.1	[1.4 ; 12.7]	15.3	[5.8 ; 24.8]	2.7	[-3.9; 9.4	
	30-44 years	0.0	ref.	0.0	ref.	0.0	ref.	
	45-59 years	-4.6	[-9.0 ; -0.3]	-9.2	[-16.1;-2.4]	-0.1	[-5.5; 5.3	
	60-75 years	-5.3	[-10.6; 0.1]	-19.0	[-27.2 ; -10.8]	9.7	[2.8;16.5	
		Pro	cessed meat					
angu	age regions							
	German	0.0	ref.	0.0	ref.	0.0	ref.	
	French	-2.9	[-5.1 ; -0.7]	-5.7	[-9.4 ; -2.1]	-0.6	[-3.4; 2.2	
	Italian	0.2	[-4.0; 4.4]	0.8	[-5.9; 7.4]	-0.5	[-5.9; 4.8	
		Red	Meat	-				
Age-g	roups							
	18-29 years	-2.8	[-6.7; 1.1]	-1.1	[-7.9; 5.8]	-3.4	[-7.9; 1.1	
	30-44 years	0.0	ref.	0.0	ref.	0.0	ref.	
	45-59 years	-1.0	[-4.0; 2.0]	-2.6	[-7.5; 2.4]	0.4	[-3.3; 4.0	
	60-75 years	-0.8	[-4.5; 2.9]	-6.3	[-12.2 ; -0.3]	5.2	[0.5; 9.8	
		Whi	te Meat					
Sex								
	Men	0.0	ref.					
	Women	-0.8	[-3.0; 1.4]					
Age-g	roups							
	18-29 years	8.4	[4.8 ; 11.9]	13.9	[7.7 ; 20.1]	4.9	[0.8; 9.1	
	30-44 years	0.0	ref.	0.0	ref.	0.0	ref.	
	45-59 years	-1.5	[-4.3; 1.2]	-3.0	[-7.4;1.5]	0.6	[-2.8; 3.9	
	60-75 years	-2.1	[-5.5; 1.3]	-5.5	[-10.8 ; -0.1]	2.2	[-2.1; 6.4	

# Further significant association p-value < 0.05

- Overweight and obesity was positively associated with consumption of all meat categories
- · Participants with children consumed more all and white meat than participants living with a partner but without children
- Tertiary education was negatively associated with all, processed and red meat consumption, than lower education levels.

## **Discussion & Conclusion**

Mean daily meat consumption of 109g consists of 37g red, 27g white, 43g processed and 2g unspecified meat.

Most prevalent differences in meatcategory consumption were between the sexes except for white meat and the linguistic regions. Swiss residents of different

- demographic groups may face different benefits & risks from their meat consumption.
- Consumption of separate meat categories should be monitored.

Energy-standardized meat consumption of the oldest age group was not significantly associated with meat consumption although the absolute meat consumption decreased with age.

Regression analysis revealed significant associations for Age, BMI, Smoking, Household status, Education & Nationality for which further study is warranted.

Results of the sex-specific regression suggest the need for separate analysis of men and women.